

STATUS AND DISTRIBUTION OF
BIODIVERSITY IN WESTERN GHATS OF
KARNATAKA

K S MURALI
RESEARCH ASSOCIATE
TATA ENERGY RESEARCH INSTITUTE
50/7, PALACE ROAD
BANGALORE 560 052

TEL : 91-80-2268296
FAX : 91-80-2265760

REPORT SUBMITTED TO
OVERSEAS DEVELOPMENT ADMINISTRATION
FORESTRY OFFICE
B2/16, VASANT VIHAR
NEW DELHI

SUMMARY

A literature survey work was carried out to understand the amount literature available on the distribution biodiversity of western ghats, to identify the priority areas for conservation efforts, to identify the taxa which need more attention and to identify the areas which were hitherto neglected and need immediate attention. Various research institutions and universities were visited for the purpose and the literature on the published work on the western ghat was compiled. The results indicate that the studies are concentrated in Uttara Kannada, Dakshina Kannada, Mysore and Kodagu districts and the districts which need more attention for the bio-diversity studies are Shimoga, Chikmagalur, Dharwar, Hassan and Belgaum. Further, our data suggest that mammals and tree species are reasonably well studied and those which further need attention are invertebrates, microbes, freshwater fishes and reptiles.

The setting:

Western ghats, richly endowed with floral, faunal and habitat diversity, is one of the twelve megadiversity centres of the world. In India, western ghats stands second only to the Himalayan tracts with respect to biological diversity. This include hill ranges from the river Tapti in the north till Kanyakumari in south for about 1600 kms (between 8° – 21° N and 73° to 78° E). The chain of hills almost runs paralelly to the western coast of India. Due to this setting western slope of the ghats receives most of the rain from south west monsoons as they face the windward direction and eastern slope forms the rain shadow region as they face the leeward direction of the monsoon. There is great variation in number of rainy months over the entire ghats, the southern parts receiving rains for nearly 8 months and the northern parts receive 4 months. Bulk of the rain received in the peninsular India is showered at western ghats which fills three major rivers of this region namely, Cauvery, Godavari and Krishna and various tributaries and streams. It's for this reason we have a varied type of habitats available in the entire range of ghats such as tropical wet evergreen, moist and dry deciduous, scrub thorny, shola-grasslands, bamboo thickets, riverine habitats, mangroves, lakes and swamps.

The Importance

A major break-through in the history of inter-continental trade is the discovery of sea route to India by Vasco-da-gama during 16th century which made many spices naturally grown in western ghats exportable to outside world, particularly Europe. Even today pepper, cardamom, cinnamomum etc., forms a bulk of Indian export market. Such a motive to encash the natural resources in a commercial sense, was started only after the visit of Vasco-da-gama which led later several Europeans to invade India for its spices. This changed the history of India, which attracted many Europeans for its spices and later became a colony of Europe. This ultimately resulted in British ruling the Indian subcontinent for nearly 180 years. It's this colonization by British lead to started exploitation of forests in a commercial manner such as extraction of timber, raw materials for various forest based industries like paper mills, match sticks and plywood making. Thus the

attraction of income generated through forest and natural resource extraction prompted Britishers to undertake extensive survey work in search of useful forests and areas for mining. This task of intense search for resource probably led to describing the vegetation of Indian subcontinent and resulted in books like many floras of India by famous taxonomists like Benthom, Hooker, Cook, Talbort, Gamble, Fyson etc., and a very good land surveying machinery was developed in India. Even today, India is one of the best surveyed and mapped country in the world.

Solely with a view to conserve the forest resources which generated lot of income to British Raj, probably for the first time in Indian history the forest areas were considered as state asset and Forest Department was constituted. Several regulations were passed restricting the entry of forest by local tribes as well as villagers and also restricting the resources harvested by local tribes and surrounding villagers for their subsistence. Even after independence, the Forest Department and its regulations continued. Probably the curbing of rights of people over forests created a rough weather among people and the forest department which led to constant clashes between these groups. Later, the focus of forest department shifted from viewing the forests just as sources of income into the non-cash benefits that the forests confer to the mankind. Therefore the legislation was brought to ban felling of trees for many kinds forestry extractions. Of late, the Government of India seems to have realized the importance of involving people in its forest conservation efforts. In the International Convention on Biological Diversity India has committed to undertake identification and monitoring of its biological diversity.

The objective

In view of signing the convention on biological diversity, it is important to take stock of biodiversity available in the western ghats. Karnataka Forest Department has long been interested in launching a programme on conservation of western ghats involving people (people's participatory project). In this connection, my attempt here is to look into the state-of-the-art of research that have been carried out in the western ghats of Karnataka. Further to understand the gaps in knowledge, particularly about the areas that are least studied, the group of organisms which are not

given proper attention and to collate the information in some organized manner. An effort was made to collect all the information available on distribution and density of various groups of organisms in the western ghats of Karnataka. For this purpose various libraries and Institutions were visited and the available information, either published in journals or books or report form, were collected. In addition, several journals were consulted to get the information. The attempt is made to collect an exhaustive list of references and wherever possible a copy of the literature was collected. Although the list of references so collected does not seem to be exhaustive, but due to limitations of time and constraints involved in travelling greater distances, many cross references were collected from the available literature source than going to the original literature source.

General remarks

A great amount of literature is (references appended as the literature on western ghats) available dealing with the biodiversity of western ghats. Unfortunately much of it is either a qualitative account describing a couple of magnificent organisms such as orchids, tigers or elephants or discovery of new genera or species or a variant morph of a species. A quantitative description of a given organism or a taxa is very limited. One of the examples of such excellent work published from Journal of Bombay Natural History Society is by Prasad et. al (1979) describing the density, distribution and abundance of mammals of western ghats of Karnataka. It would be a good step now to compare that data with the current data to understand change in population. The same procedure can be adopted to draw distribution and density maps of other vertebrates such as birds, amphibians, reptiles etc., using the scattered data available at different protected areas in the western ghats of Karnataka. Infact, for most protected areas birds data are available (although not the data on density) which can be used to draw distribution maps of various bird species.

Similarly, vegetation maps prepared by French Institute, Pondicherry, (Pascal 1986) are very useful at gross level to understand different vegetation types over the western ghats of Karnataka. This book let also gives the species available over different forest types. However, the data from the working plan of each division contains the densities of different species of trees at certain intervals

of time which can be used to classify maps based on dominant species and their densities and can draw distribution of species. Further, it can be used to understand change in species due to forestry operations and can relate the rate of change of species on the intensity of felling etc. The detailed data on distribution of only commercially important species such as Sandal (*Santalum album*), Teak (*Tectona grandis*) and Rosewood (*Dalbergia* sp) are available. Much literature from forestry journals deal with provenance trials of exotic species such as *Eucalyptus* and *Acacia equisitifolia*'s are available, whereas some of the equally important indigenous species such as *Terminalia* have not been studied. Similarly many of fruit yielding tree species which are important in their economic value to local villagers such as Jack fruit (*Artocarpus integrifolia*), Nelli (*Emblia officinalis*), Mango (*Mangifera indica*), Bore (*Ziziphus mauritiana*), Nerele (*Syzizium cumini*) have not been well studied with respect to their occurrence in density. distribution etc.

Gaps in knowledge

Among the literature available a tally was done as to how many papers deal with birds, mammals, vegetation etc., and how many of the papers deal with different districts wherein western ghat hill ranges are found and the data are given in table 1. It is implicit from the table that the districts Chikmagalur, Belgaum, Dharwar, Hassan and Shimoga have hardly any information on many taxa. Although the total number of papers available in each taxa are reasonable enough to describe the minimum literature on the concerned biota, but some districts like Uttara Kannada, Dakshina Kannada and Mysore weight more than the rest. The study are not distributed over all the districts uniformly but are confined to Uttara Kannada, Dakshina Kannada, Mysore and Kodagu. Many times, it so happened that the researchers have described their studies as western ghats, and not specified which locality within the western ghat they have worked. This data thus suggest that we have to concentrate on the areas which have not been fully explored. The districts which need to be given more attention are Chikmagalur, Shimoga, Dharwad, Hassan and Belgaum although these districts have reasonably good forest area (table 2).

The vegetation description of various areas are available in forest working plan of the

respective divisions or through the floras of the districts. However a quantitative account of vegetation are not available apart from the working plans. Emphasis should now be laid to get the quantitative data of various national parks, sanctuaries and reserved forests for the purpose of identification of Zone I area. Further, a common procedure should be followed to cover the diversity of all the strata of vegetation such as canopy, understory, shrub and herbs. For this purpose linear transects of one km long and 10 m wide can be suggested which also takes care of spatial heterogeneity. In order to have commonality the procedures followed by the team from Centre for Ecological Sciences (at Sirsi and Kumta) or by the team by French Institute (at Kodagu) or the procedures followed by the team at Tata Energy Research Institute (at BR Hills) can be made use of. Much emphasis is given to study the density of some particular timber yielding tree species. But now we should also emphasize the study of herbs and shrubs apart from studying the other less studied tree species and knowing their distribution.

Much studied species among animals are mammals and birds. Again these studied have concentrated in Uttara Kannada and Mysore. The attempts should be made to study the same in other national parks and sanctuaries in other districts also. From among the vertebrates which has been neglected are reptiles. Although there is one book by Daniels (1974) which only describes the south Indian reptiles qualitatively. No study yet been done regarding behaviour, density and distribution of these species. Much of the study of fishes have been conducted at Kali estuary and around Karwar rivers. The distribution of marine fishes and their distribution of western coast is well documented. However, much needs to be done on inland fish species of riverine habitats (fresh water fishes) and how their distribution are determined. Study on distribution of amphibians have been done to some extent by Ranjit Daniels (1992).

One of the significant taxa of animals which has been neglected are the study of invertebrates which is considered to be contributing a lot to diversity and in terms of their contribution to the total biomass. It is considered that phylum Arthropoda alone contributes greatly to the total biomass of the world. A great deal of interest need to be created among research

workers to undertake this work. The study of molluscs and other invertebrates are worth attempting as there may be interesting patterns discovered in this part of the world. We have excellent opportunity to study these systems as we have varied ecosystems in a very limited area of this western ghats. Another herculean task for biologists is to account for the diversity of lower organisms such as fungi, bacteria, mosses and lichens. As one realizes the tasks ahead to study even one group of insect itself is difficult, it is more difficult to take up diversity studies on these lower organisms. I would not pretend to recommend any suggestions at this stage. A lot of discussion is necessary before a criteria to study lower organisms are attempted.

In total, although it appears a large body of literature is available, we will have to scrutinize many papers for their worth in considering the work as done, before taking up planning work on the distribution of biodiversity of western ghats. I feel much more exercise is needed to compile already published information and take up further survey work at different districts. It may also be necessary to process published information for the purpose and then look for strategies to plan further.

Acknowledgements : I thank Dr. Michael Harrison, Consultant to JDA, Mr. Sadashivaiah, Chief conservator of forests (Western ghats development) for providing a lot of interaction regarding this project. I thank Mr. H V Dayal, Senior fellow, TERI, Bangalore, for encouragement and Mr. Srinivasa Rao, Mr. V T. John (from UAS) and Mr. Ravi Hegde (From TERI) in collection of literature.

Table 1 : The distribution of scientific studies over different districts of western ghats and over different plant and animal groups.

District	General	Tree	Herb	Shrub	Orchid	Mammal	Fish	Bird	Amphi- bians	Microbe	Wildlife general	Insect	Ecology & Environment	Total
Uttara Kannada	6	1					2	3		4	1	3	2	22
Dakshina Kannada							7		1	5		2		15
Shimoga				1				2						3
Dharwar	1					1		2						4
Chikmagalur														0
Hassan	1													1
Mysore	2					3		2			1		2	10
Kodagu	4			1										5
Belgaum														0
Western Ghats (general)	16	18	5	4	2	10	4	4	3		9	3	4	63
Total	31	19	5	6	2	14	13	13	4	9	11	8	8	140

Table 2. The geographical area and forest area (in km²) and per cent forested area in different districts of western ghats in Karnataka. (Source: Karnataka State Forest Annual report)

District	Geographical area	Forest area	% forest area
Belgaum	13444	2245.67	16.70
Chikmagalur	7221	2179.08	30.18
Dakshina Kannada	8336	5182.30	62.17
Dharwad	13782	1436.73	10.42
Hassan	6622	541.07	8.17
Kodagu	4108	1259.52	30.66
Mysore	12463	4129.62	33.14
Shimoga	10576	3270.16	30.92
Uttara Kannada	10247	8291.51	80.92

Literature on Western Ghats

- Achutan Kutty, C.T. & Nair, S R.S. 1982. Preliminary account on the distribution of decapod larvae in Konkan waters. *Mahasagar* 14(3): 211-214
- Achutan Kutty, C.T., Nair, S.R.S., Devassy, V.P & Nair. V.R. 1981. Plankton composition in two estuaries of the Konkan coast during pre-monsoon season *Mahasagar* 14(1) : 55-60
- Adkoli, N.S. 1976. Western ghats of Karnataka *Myforest* 12(4): 161-174
- Akbar Shah, A. 1990 Basal area distribution in tropical rainforests of Western Ghats *Indian For.* : 356-368
- Alva, U.T. 1990. Wildlife in Karnataka. *Myforest*, 25(3): 299-303
- Anonymous. 1964 Sanctuaries in Mysore state *Myforest* 1(1):42-49
- Ansari, Z.A. 1978 Meiobenthos from the Karwar region (central west coast of India). *Mahasagar* 11 : 163-167
- Appayya, M.K. 1991. Wildlife in Karnataka - A report for 1990-91. *Myforest* 27 : 309-312
- Appayya, M.K. 1985. Sea turtle conservation with particular reference to Karnataka. *Myforest* 21(2) : 89-90
- Appayya, M.K. 1993. Elephants in Karnataka (India) - A status report *Myforest* 29 :165-170
- Arora, R.K. 1950. The flora of North Kanara. *Indian For.* 86(10): 609-616
- Arora, R.K. 1960. The botany of Coorg forests. *Proc. Natl. Acad. Sci.* 30(b): 289-305
- Arora, R.K. 1964. The botany of Coorg forests - II. *Proc. Natl. Acad. Sci.* 38(b): 100-112
- Arora, R.K. 1965 Ecological notes on vegetation of coorg district, Western Ghats. *Indian For.* 91(10): 722-742.
- Babu, K.S. & Neelakantan, P. 1982. Biology of *Liza parsi* in Kali estuary, Karwar. *Mahasagar* 16(3): 381-389
- Balaiah, B 1975 Elephants in Karnataka. *Myforest* 11(1):81-84
- Banasode, 1948 From scrub forest to shrub forests of Bijapur district and the problem of afforestation. *Indian For.* 12: 405-408.
- Barnes, E 1944. Notes on the flowering plants of Biligiri Rangan Hills. *J. Bombay Natl. Hist. Soc.* 44: 436-459
- Basappanavar, C.H. 1985. Saving the tiger in Bandipur National Park. *Myforest* 21(3): 171-174
- Basappanavar, C H. 1985 Twelve years of project tiger in Bandipur National Park. *Project tiger KFD.* 1-14

- Basavalingappa, S. 1981. Foraging behaviour of the termite *Odontotermus wallonensis* (Isoptera:Termitidae). G K Veeresh Progress in Soil biology and ecology in India. 37: 272
- Belliappa, K.A. 1987. Logging in Karnataka. KFD Seminar on Logging development. pp 36-37
- Bengeri, K.N. 1979. Virgin game sanctuary. Myforest 15(3): 187-188.
- Benkappa, S. Al. 1978. Influences of some oceanographic conditions on the fishes of Mukka kamp along South Canara coast. Indian J Mar. Sci. 8: 235-282
- Benkappa, S. Al. 1979. Distribution of Plankton in the Arabian sea between Mukka and Kali along South Kanara coast. Mysore J. Agric. Sci. 13: 454-463
- Bhagawat, S.R. 1975. Our vanishing forests. Myforest 11: 123-126.
- Bharati, S.G. et al. 1980. Limnological studies in ponds and lakes of Dharwar. Comparative phytoplankton ecology of 4 water bodies. Phykos 19(1): 27-43
- Bhaskar, V. 1982. Karnatakadalli gandada marada abhivruddhi. Myforest 2:67-68.
- Bhat, D.M. 1992. Phenology of tree species of tropical moist forest of Uttara Kannada. Karnataka. J. Biosci. 17(3): 325-352
- Bhat, D.M. et al. 1986. Biological species diversity in some localities of western ghats. Tech Rep No. 33 CES, IISc., Bangalore pp 1-25
- Bhat, D.M. 1987. Plant diversity studies in Uttara Kannada. Tech rep No.9, CES, IISc., Bangalore.
- Bhat U.G. & al 1988. Environmental characteristics of the mine and estuarine habitats of Karwar - An overview. J. Indian Fish. Assoc. 17: 401-412
- Bhat, U.G. & Neelakantan, B. 1983. Some aspects of the fishery of fringe scale *Sardine sardinella fibriata* in Uttara Kannada, Karnataka. J. Karnatak Univ. Sci. 28: 142-147
- Bhat, U.G. & Neelakantan, B. 1985. Edible clam resources of Kali estuary, Karwar. Sea Food Export J. 17(2): 21-25
- Bhat, U.G. & Neelakantan, B. 1985. Composition and distribution of benthos in Kali estuary, Karwar. J. Indian Fish Assoc. 14 & 15: 27-35
- Bhat, U.G. & Neelakantan, B. 1991. Distribution of meio benthos in relation to environmental parameters in the Kali estuary, Karwar. Comp. Physiol. Ecol. 16(2): 60-68
- Bhat, B.V. & Gupta, T.R.C. 1983. Zooplankton distribution in Nethravathi-Gurupur estuary, Mangalore. Indian J. Mar. Sci. 12: 36-42
- Bhat, H. R., Srinivasan, M. A. 1978. Recent sightings of *Macaca sinensis* at Mastimane ghat North Kanara district, Karnataka. Indian J. For. 75(2): 476.
- Bhat, K.G. 1985. Eco-development of western ghats - role of wood processing. Indian. J. For. 8: 102-108

- Bhat, K.M., Rajan, B.K.C. & Jain, J.C. 1979. Mahagony (*Sweitenia mahogany* Jacq.) in Karnataka. *Myforest* 3: 41-43
- Birasal, N.R. et al. 1988. Ecology of Zooplankton of supra reservoir of Kali river during filling phase (western ghats, Karnataka). *J. Freshwater biol.* 1(1): 15-28
- Boopaiah, B.A. & Neelakantan, B., 1982. Ecology of a tidal pond in an estuarine environment in Karnataka. *Mahasagar* 15: 29-36
- Chinnamani, S. 1968. Grasslands of Bellary black cotton soils. *Indian For.* 94(3): 225-229
- Chinnamani, S. 1982. Grasslands of South India and its ecology, floristic composition and utilization. *Bull. Bot. Surv. India.* 23 : 128-134
- Cleghorn, H. 1861. Forests and gardens of South India. W. H Allen and Co., London. pp 412
- Daniels, J.C. 1963. Field guide to the amphibians of western India. Part I & II. *J. Bombay Natl. Hist. Soc.* 60: 415-438; 690-702
- Daniels, J.C. 1975. Field guide to the amphibians of western India. Part III. *J. Bombay Natl. Hist. Soc.* 72: 506-522
- Daniels, J.C. 1975. The book on Indian reptiles. *Bombay Nat. Hist. Soc.*
- Dasappa, 1989. Geographical distribution problems of regeneration (natural and artificial) in teak *Tectona grandis*. *Myforest* 25: 337-355
- David, A. et. al. 1969. Limnology and fisheries of the Tungabhadra reservoir. *Bull. Central Inland fisheries Res. Inst.* 13.
- Davidson, J. 1898. Birds of the North Canara - I & II. *J. Bombay Natl. Hist. Soc.* 11:652-679
- Deshmukh, D.K. 1975. Regeneration of rose wood. *Myforest* 10: 87-93
- Devassy, V.P. & Groes, J.I. 1989. Seasonal patterns of phytoplankton biomass and productivity in a tropical estuarine complex (west coast, India). *Proc. Indian Aca. Sci. (Plant Scie)* 99(5): 485-501
- Devi Prasad, A.G. 1988. Distribution and economic potential of *Dalbergia* in Karnataka. *Myforest* 4: 241-249
- Dharmaraj, K. & Nair, N.B. 1981. Timber boring animals of Talapady mangroves in adjoining areas and with a record of boring into living trees. *Mahasagar* 14 : 159-162
- Divakar, K.M. 1986. A study on the mangroves of the Kali and Aghanashini river estuaries of Uttara Kannada. *J. Indian Bot. Soc.* 65: 111-116
- Gadagkar, R. Chandrashekara, K., & Nair, P. 1990. Insect species diversity in tropics : sampling methods and a case study. *J. Bombay Natl. Hist. Soc.* 87(3): 337-353
- Gadagkar, R. Nair, P. Chandrashekara, K. Bhat, D.M. 1991. Ant species richness and diversity in some selected localities in the western ghats CES, IISc., pp 10

- Gadgil, M. 1979. Hills, dams and forests, some field observations from the western ghats. Proc. Indian Acad. Sci. 2(3): 291-203
- Gadgil, M. 1983. Status of wild mammals in Karnataka. In (ed) C. J. Saldanha. Karnataka State Environment Report. pp 22-35
- Gadgil, M. 1987. Geographical distribution of Plant species in Karnataka. Tech Rep. 32. CES, IISc., Bagalore
- Gadgil, M. & Meher Homji, V.M. 1990. Ecological diversity. In Daniels, J.C. Serrao, J.S. (ed) Conservation in developing countries - Problems and prospects. Bombay Nat. Hist. Soc pp 175-198
- Gadgil, M. & Meher Homji, V.M. 1991. Status survey of vegetation types of India (western ghats). CES, IISc., Bangalore, pp 40
- Ganapati, K.T. 1976. Project Tiger. Myforest 1(2): 51-59
- Gaonkar, V.S., Vartak, V.D. & Kumbhajkar, M.S. 1990 Indigenous fuel-wood species from western ghats of Maharastra. Indian For. 3: 233-237
- George, P.C. 1900. Observation of the crab fishery of Mangalore coast. Indian J. Fish. 43-52
- Ghorpade, K. D. 1973. Priliminary note on ornithology of Sandur, Karnataka. J. Bombay Nat. Hist. Soc. 70(3): 499-531.
- Gopal Reddy, K. & Chandrashekara Gupta, R.T. 1988. Studies on the phytoplankton of Mulki estuary, Dakshina Kannada. Mysore J. Agric. Sci. 22:258-265
- Govindakrishnan, P. M., Verghese, A. and Chakravarthy, A. K. 1978. Occurrence of red headed murline in Bangalore, Karnataka. J. Bombay Nat. Hist. Soc. 75(2): 487.
- Green, S. & Minkowski, K. 1977. The lion tailed monkey and its south Indian rainforest habitats. Primate conservation. New York pp 289-337
- Gunaga, Vishnudas, Neelakantan, K. & Neelakantan, B. 1990. Tidal and lunar influences on penacid prawn seed abundance in the Kali estuary, Karwar. Mahasagar 23: 63-69
- Gurudeva, M. R., Balakrishna Gowda, Rajanna, M. D. 1981. *Ophioglossum costatum* R. Ber. new to Karnataka. Indian J. For. 4(4) 326.
- Hegde, G.R. 1985. On the succession of algae in a temple tank at Darwar, Karnataka. Geobios 12(6): 261-263
- Hussain, K.M. 1971. The ruthless India wild dog or Dhole (*Cuon alpinus* Pallas). Myforest 8(3): 1-5
- Hussain, K.M. 1975. Flora and fauna of Karnataka part I. Myforest 10(3): 11-19
- Hussain, K.M. 1975. FLora and fauna of Karnataka Part II. Myforest 10(4): 3-6
- Hussain, K.M. 1975. Flora and Fauna of Karnataka, part III. Myforest 11(1): 3-10
- Hussain, K.M. 1975. Flora and Fauna of Karnataka, part IV. Myforest 11(2) 55-62, 105-114

- Jagadeesh, K.S & Geetha, G.S. 1989. Utilization of *Eupatorium*—the micrological approach. *Myforest* 25: 125–129
- Jain, S.K. 1939. The montane evergreen forests, Bisle region. *Indian For* 65: 182–201
- Jain, S.K. 1944. The ghat rain forests, Agumbe–Kilandur zone. *Indian Forester* 67(4): 184–203
- Jain, S.K. 1950. Evergreen montane forests of western ghats of Hassan district, Mysore. *Indian For.* 76: 18–30; 69–82; 121–132
- Jayaraj, E.G. & Reddy, M.P.M. 1984. Possible effects of bottom water currents of hydrographic features on demersal fisheries off Mangalore, South Kanara. *Malasagar* 17: 1–7
- Johnsingh, A. J. T. 1983. Large mammalian predators in Bandipur. *J. Bombay Nat. Hist. Soc.*, 80(1): 1–58.
- Kadambi, K. 1941. The evergreen ghat rainforests, Agumbe–Kilandur. *Indian. For.* 67: 184–203
- Kadambi, K. 1942. The evergreen ghat rainforests of the Tunga and Bhadra river sources. *Indian For.* 68: 233–240
- Kadambi, K. 1949. On ecology and silviculture of *Dendrocalamus strictus* in the bamboo forests of Bhadravathi division, Mysore state. *Indian For.* 75(9): 289–299
- Kadambi, K. 1950. Evergreen, montane forests of the western ghats of Hassan district, Mysore state. *Indian For.* 76: 18–30; 69–82; 121–132
- Kaikini, N.S & Shyamsundar, S. 1967. Oil palm in Mysore forests. *Myforest* 4(1): 1–6
- Karanth. U.K. 1982. Bhadra wildlife sanctuary and its endangered ecosystem. *J. Bombay. Nat. Hist. Soc.* 79(sup): 79–86
- Karanth. U.K. 1985. Indian black crested buzzard; a sighting record from Karnataka. *J. Bombay Natl. Hist. Soc.* 82(1): 183.
- Karanth. U.K. 1984. Conservation plan for the lion tailed macaque and its rainforest habitats in Karnataka DEE, GOK, pp 118
- Karanth. U.K. 1986. A possible sighting record of malabar civet (*Viverra zibethica* Blyth) from Karnataka. *J. Bombay Natl. Hist. Soc.* 83(1): 192
- Karanth, U.K. 1988. Analysis of predator–prey relations in Bandipur tiger reserve with reference to census reports. *J. Bombay Nat. Hist. Soc.* 85(1): 1–8
- Kelanjer, A.K. 1976. A preliminary survey of the orchids of Coorg district. *J. Mysore Univ.* 27: 189–196
- Keshava Murthy, K.R. 1982. *Medico-Botany of Karnataka - II* 2:43–58
- Keshava Murthy, K.R. & Yoganarasimha, S.N. 1990. *Flora of Coorg* Vimsa: publ. Co.,
- Krishnappa, H.P. 1982. Improvement fellings and regeneration of hardwood in Deciduous forests – Sargod method. *Myforest* 18(2): 97–101

- Kulkarni, C. V. and Ogha, C. V. 1978. The present status of mahseer fish and artificial propagation for Khudrea (Sykes)
- Kushalappa, K.A. 1988 Comparative biomass of *Acacia auriculiformis* and *Leucana leucocephala* trees from moist region of Karnataka. Myforest 24(1): 12-15
- Kushalappa, K.A. & Akbar Shah. 1978. *Gmelina arborea* Roxb. - Provenance trials in Karnataka. Myforest 14(2): 87-91
- Kushalappa, K.S. 1989. Biomass studies in *Callindra* in Karnataka. Myforest 25(4): 325-329
- Kusuma, M.S., Neelakantan, B, & Konnur, R.G. 1988. Plankton distribution in the Kali estuary, Karwar, Central west coast of India. Environ. Ecol. 6: 115-119
- Lakshmana, A.C. 1983. Orchid of western ghat. Myforest 24(3): 201-202
- Lakshmana, A.C. Vedant, C.S. & Jaya, M.M. 1992. Status report on canes (*Calamus rotang*, *C. thwaitesii*, and *C. vattayila*) in Kodagu district of Karnataka. Myforest 28(4): 343-347
- Larsen, T.B. 1987. The butterflies of Nilgiri mountains of South India. (Lepidoptera:Rhopalocera).
- Lott, E.J. 1985. European bee eaters in Karnataka. J. Bombay Natl. Hist. Soc. 82(2): 411
- Madhyastha, A. N. 1987. First report of masked booby, *Sula dactylata* from the shores of Karnataka.
- Madhyastha, M.N. Sharath, B.K. & Rao, I.J. 1986 Preliminary studies on marine turtle hatchery at Bengre beach, Mangalore. Mahasagar 19(2): 137-140
- Mallikarjunaiah, T.S. 1978. Some of our fascinating birds Part I Myforest 14(2): 97-101
- Mallikarjunaiah, T.S. 1979. Some of our fascinating birds part II Myforest 15(2): 91-95
- Mannikeri, M.S. et al. 1991. A note on ecology and distribution of recent freshwater ostracodes from Alnavar, North Kanara, Seminar on Ecology of Western ghats. Karnatak Univ. pp 10
- Meher Homji, V M. 1974. On the origin of the tropical evergreen forests of South India. Int. J. Ecol. Envir. Sci. 1: 19-39
- Mohammad Hussain. A. M. 1962. Yield regulation in the sandal forests of Madras state. Indian For. 88(8): 543-595.
- Muddanna, V. 1971. Contributions to the knowledge of aquatic weeds in fisheries tanks of Mysore. their distribution and influences on fisheries. Tech. Report UAS, Bangalore
- Nadakarni, V.B. et al. 1988. A survey studies on biology of endangered species of amphibians and reptiles of western ghats- Uttara Kannada, and Chikmagalore. Karnataka western ghats development programme Karnatak Univ. pp 40
- Nagarajaiah, C.S. 1983. Studies on hydrobiology of brackish water ponds of Nethravathi estuary, Mangalore, Dakshina Kannada. Mysore J. Agric. Sci 17: 212

- Nair, K.K.N. 1986. Additions to Gamble's flora of Presidency of Madras (1915-1935) from the state of Kerala, Tamilnadu, Karnataka and Andhra Pradesh upto 1982 Indian J. For. 9(3): 204-219
- Nair, N.C., Chandrabose, M. and Srinivasn, S. R. 1980. A further contribution to the weed flora of South India. Indian J. For. 3(2): 111-115.
- Nair, N.C., Chandrabose M. & Srinivasan, S.R. 1980. A further contribution to the weed flora of South India Indian J. For. 3(1): 56-59.
- Nair, N.C. & Daniel, P. 1980. The floristic diversity of the western ghats and its conservation : A review. Proc. Indian Acad. Sci (Animal and Plant Sci) Suppl. 127-163
- Nair, P.V. & Gadgil, M. 1977. The status report of elephant populations in North Kanara. Tech. Rep. no 4 IISc., Bangalore-12
- Nair, P.V. & Gadgil, M. 1980. The status and distribution of elephant populations in Karnataka. J. Bombay Natl. Hist. Soc. 75(suppl): 1000-1016
- Naithani, B.D. 1966. Studies on the flora of Bandipur reserve forest, Mysore state. Bull. Bot Surv. India. 8: 253-263
- Nanayya, K. M. 1949. Sandal forests of Coorg. Indian For. 3:87-90.
- Nanayya, K. M. 1975. Regeneration of wild date palm (*Phoenix sylvestris*, Roxb.). Myforest 26(1): 118-120.
- Nanjundappa, D. 1957. A note on *Eucalyptus* plantations in old Mysore state. Indian For. 4: 280-286
- Nayak, V. N. and Kakate, V. S. 1978. Occurrence of the hermit crab *Dandranus setifer* at Karwar.
- Nayak, V.N. & Neelakantan, B. 1989. A new species of hermit crab *Diogenes karwarensis* from the west coast of India. J. Bombay Natl. Hist. Soc 86(1): 71-77
- Neelakantan, B. 1992. Aquaculture studies along the fringes of mangrove habitats of the UttaraKannada districts of Karnataka. Dept of Marine Biology Karnataka Univ Karwar. pp 12-27
- Neelakantan, B 1985. On the molluscan resources in Uttara Kannada. Fisheries Tech. pp 88-90
- Neelakantan, B., Nauk, U. G., Prasad, P. N. and Kusuma, M. S. 1988. Hydro-biology of Kali estuary and adjoining brackish water systems, Karwar. Environment and Ecology 6(1): 139-147.
- Neginhal, S. G. 1971. Birds of Dharwar district. Myforest 3(3): 23-24.
- Neginhal, S. G. 1974. Blackbuck of Ranibennur. Myforest 10(2): 13-16.
- Neginhal, S. G. 1978. Tigers. Myforest 14: 167-168.
- Neginhal, S.G. 1980. Ecological impact of afforestation at the Ranibennur black buck sanctuary. J. Bombay Natl. Hist. Soc. 75(supl) 1254-1258

- Neginhal, S. G. 1982. Ecology and behaviour of the Great Indian Bustard (*Choriatris nigriceps*). Myforest, 2:59-66.
- Neginhal, S. G. 1986. Some beautiful trees of Bangalore. Myforest 22(4): 241-244.
- Neginhal, S. G. 1993. The project tiger. Myforest 29(3): 161-164.
- Neginhal, S. G. 1993. Wildlife through history in Karnataka. Myforest 29(3): 215-224.
- Neginhal, S. G. 1982. The birds of Ranganatittu. J. Bombay Natl. Hist. Soc. 79(3): 581-593
- Neginhall, S.G. 1991. Bandipur ecosystem then and now (in 1972 and in 1991). Myforest 27(3): 235-238.
- Pai, R. & Reddy, M.P.M. 1981. Influence of some oceanographic factors on the trawl catches off Malpe, south Karnataka. mss
- Pascal, J.P. 1986. Forest map of South India. French Institute, Pondicherry.
- Pascal, J.P. & Ramesh, B.R. 1987. A field key to the trees and lianas of the evergreen forests of the western ghats (India). French Institute, Pondicherry. pp 1-236
- Prabhu, M. S. et al. 1974. Fisheries resources of Ullal (Mangalore) in relation to certain environmental factors during 1963-1967. J. Bombay Nat. Hist. Soc. 71(1): 95-98.
- Prabhu, M.S. & Dhulkhed, M.H. On two new varieties of Indian Oil Sardine, *Sardinella longiceps* Val. Mahasagar 5(1): 27-30
- Prabhu, V. & Reddy M.P.M. 1987. Macrobenthos & sediment distribution in relation to demersal fish catches off Baikampad-Suratkal, South Kanara coast. Indian J Mar Sci 16 60-64.
- Prabhu, V. & Reddy, M.P.M. 1987. Distribution of Nutrients and Plankton in the Nearshore waters off Baikampady-Suratkal. South Kanara Environ. Ecology 5: 247-252
- Prajapathi, R.C. 1989. Depletion of forest wealth of western ghats. Myforest 25(4): 361-364
- Prakash, H. S. & Sreeram Reddy, G. 1984. Distribution of *Drosophila* species and their diversities in the tropical rain forests of Western Ghats. J. Bombay Nat. Hist. Soc. 81(1). 323-345
- Prasad, S.N., Nair, V P., Sharatchandra, H.C. & Gadgil, M. 1978. Factors governing the distribution of mammals in Karnataka J. Bombay Nat. Hist. Soc. 75(3): 718-743.
- Prasad, S.N. & Sharatchandra, H.C. 1984. Primary production and consumption in the deciduous forest ecosystem of Bandipur in South India. Proc. Indian Acad. Sci (Plant Sci) 93. 83-97
- Prasad, S.N. & Hegde, M. 1986. Phenology and seasonality in the tropical dry deciduous forest of Bandipur, South India. Proc. Indian Aca. Sci (Plant Sci). 96(2): 121-133
- Prasad, P. N., Sudarshan, R. and Neelakantan, B. 1988. Feeding ecology of mud crab *Scyella serrata* in Sunkery back waters. Karwar J. Bombay Nat. Hist. Soc. 85(1): 200
- Puttaiah, E.T. & Somashekar, S.K. 1987. On the phenology of demersals in the lakes of Mysore district. Geo bios new reports 6: 132-137

- Quereshi, T.M. 1983. A note on distribution, geology, etc., of sandalwood in Bombay state. Indian For. 318-321.
- Ragunath, V., Subramanya, S., Shyamal, Lokesh, R. and Vasudeva, R. 1992. A preliminary survey of Gudavi bird sanctuary. Myforest 28(3): 265-274.
- Rahman, M.F. & Rajagopal, K.V. 1978. Occurrence of *Ichthyophis beddomei* Peters in South Kanara, Karnataka, Science and Culture. pp 187-188
- Rai, S.N. 1979. Rate of growth of *Hopea parviflora* Bedd. Myforest 1: 31-39.
- Rai, S.N. 1980. Floristic composition and survival patterns of tropical rainforest tree species of western ghats, India. Myforest 17: 101-110
- Rai, S. N. 1981. Regional water table for Indian copal (*Vateria indica* Linn) tree and its certain other relationships (data from Karnataka). Indian J. For. 4(8): 99-101
- Rai, S N. 1990 Restoration of degraded tropical rainforests of Western ghats. Indian For. 116(3): 179-188
- Rai, S.N. & Proctor, J. 1986. Ecological studies on four rainforests in Karnataka. J. Ecol. 74(2): 455-463
- Rajagopal, D. 1982. Relative indices of termites on exotic species of *Eucalyptus* in Karnataka. Myforest 18(1): 9-13
- Rajagopal. K.V., Mohan Joseph, M., & James, P.S.B.R. 1978. A list of fishes of Karnataka. UAS Tech Rep Series. 18
- Rajagopala Shetty, K. 1991. Gudvi - The picturesque bird sanctuary. Myforest 27(3): 241-243
- Rajagopalam, P.K. 1972 *Ixodid* ticks (Acarina :Ixodidae) parasitizing wild birds in the Kyasanur forest disease areas of Shimoga district, Mysore district, Karnataka. J. Bombay Nat. Hist. Soc. 69(1): 55-78.
- Rajeev, B.M T. 1993 Eco-development - An unique approach in Bandipura National Park. Myforest 29(3). 157-195
- Ramakrishna, A C & Lakshmana, A.C. 1974 Forest of Karnataka and the five year plans. Myforest 10(2) 5-11
- Ramaswamy, M.N., Range gowda, D., & Krishnamoorti, R. *Oscimum kilimanjaricum* in Mysore state Indian For 612-619
- Ramu, G. Check-list of butterflies of lower droog-nellithurai forest, western ghats. 12(3): 238-240
- Ranjit Daniels, R J 1990. Changes in bird fauna of Uttara Kannada, India in relation to changes in landuse over the past century Biol Conserv. 52: 37-48
- Ranjit Daniels, R J. 1991. The problem of conserving amphibians in the western ghats, India. Curr Sci 60(11) 63-632

- Quereshi, T.M. 1983. A note on distribution, geology, etc., of sandalwood in Bombay state Indian For. 318-321.
- Ragunath, V., Subramanya, S., Shyamal, Lokesh, R. and Vasudeva, R. 1992. A preliminary survey of Gudavi bird sanctuary. Myforest 28(3): 265-274.
- Rahman, M.F. & Rajagopal, K.V. 1978. Occurrence of *Ichthyophis beddomei* Peters in South Kanara, Karnataka, Science and Culture. pp 187-188
- Rai, S. N. 1979. Rate of growth of *Hopea parviflora* Bedd. Myforest 1: 31-39.
- Rai, S.N. 1980. Floristic composition and survival patterns of tropical rainforest tree species of western ghats, India. Myforest 17: 101-110
- Rai, S. N. 1981. Regional water table for Indian copal (*Vateria indica* Linn) tree and its certain other relationships (data from Karnataka). Indian J. For. 4(8). 99-101
- Rai, S.N. 1990. Restoration of degraded tropical rainforests of Western ghats. Indian For. 116(3): 179-188
- Rai, S.N. & Proctor, J. 1986. Ecological studies on four rainforests in Karnataka J. Ecol. 74(2): 455-463
- Rajagopal, D. 1982. Relative indices of termites on exotic species of *Eucalyptus* in Karnataka. Myforest 18(1): 9-13
- Rajagopal, K.V., Mohan Joseph, M., & James, P.S.B.R. 1978. A list of fishes of Karnataka. UAS Tech Rep Series. 18
- Rajagopala Shetty, K. 1991. Gudvi - The picturesque bird sanctuary. Myforest 27(3): 241-243
- Rajagopalam, P.K. 1972. *Ixodid* ticks (Acarina :Ixodidae) parasitizing wild birds in the Kyasanur forest disease areas of Shimoga district, Mysore district, Karnataka. J. Bombay Nat. Hist. Soc. 69(1): 55-78.
- Rajeev, B.M.T. 1993. Eco-development - An unique approach in Bandipura National Park. Myforest 29(3): 187-195
- Ramakrishna, A.C. & Lakshmana, A.C. 1974. Forest of Karnataka and the five year plans. Myforest 10(2): 5-11
- Ramaswamy, M.N., Range gowda, D., & Krishnamoorti, R. *Oscium kinnayanicum* in Mysore state. Indian For. 612-619
- Ramu, G. Check-list of butterflies of lower droog-nellithurai forest, western ghats 12(3): 238-240
- Ranjit Daniels, R.J. 1990. Changes in bird fauna of Uttara Kannada, India in relation to changes in landuse over the past century. Biol Conserv. 52: 37-48
- Ranjit Daniels, R.J. 1991. The problem of conserving amphibians in the western ghats. India. Curr. Sci 60(11): 63-632

- Sampathkumar, R. 1987. Distribution of Plankton of waters off Mangalore. Mysore J. Agric. Sci. 21: 64
- Satyanarayana, Y. 1958. Ecological studies of the evergreen vegetation of western ghats. Symp. On Humid tropics vegetation. UNESCO pp 196-211
- Sharatchandra, H.C. & Gadgil, M. 1950 On the time budget of different life history stages of chital (*Axis axis*). J. Bombay Nat Hist Soc. 75(suppl): 949-960
- Sharathchandra, H. C. and Gadgil, M. 1955. A year of Bandipur. J. Bombay Natl. Hist. Soc. 72(3): 623-647
- Shastri, S S. 1978 Trends in marine fish landings in India. Mahasagar 11(12): 73-81
- Shivabalan, A. et al. 1990 Mangroves of Uttarakannada, Karnataka : Present status and management Environ. Ecol. 9(2): 441-444
- Shivaramaiah, S K. 1979. Brief note on elephants. Myforest 45-50
- Singh, A. K and Lala, 1983. Observauon on the movement of two captive reared mugger crocodiles, *Crocodylus palustris*. Lessons when returned to the wild. J. Bombay Nat. Hist. Soc. 80(1): 86-90.
- Singh, F. 1988. Exquisite orchids from western ghats (India) *Dendrobium lawianum*. Myforest 24(3): 169-171
- Somashekar, R K. 1982. Algal flora of river Cauvery, Cyanophyceae and Chlorophyceae. Phykos 22: 73-80
- Somashekar, R.K. 1983. A study on the microflora of river Cauvery. Sci. Culture 49: 135-137
- Somashekar, R K. 1984 Contribution to the algal flora of river Kapila I. Cyanophyceae and Chlorophyceae. Phykos 23 116-124
- Somashekar, R.K. 1984 Contribution to the algal flora of river Kapila II Diatoms. Phyko 23: 125-129
- Somashekar, R.K. et al 1989 Bacteriological quality of Cauvery waters - A preliminary analysis. Indian J. Environ health. 26: 264-268
- Sridhar, K.R. & Kaveriappa, K M. 1991 A note on marine fungi from Mangalore coast. Mahasagar 24(1): 63-66
- Srinathi, R. A.Kulkarni, H D and Venkateshan, K. R. 1980 Selection of sandal for spike resistance and other qualities Indian J For 14: 303-305.
- Srinivasan, M. A. 1973. Trapping of small in relation to the vegetation types in Kyasanur forest disease area, Mysore State. India J. Bombay Nat. Hist. Soc. 70(3): 488-492.
- Subramanian, K.N. 1986. Eco-restoration of the high altitude forest of western ghats area of Tamilnadu-An urgent need for their eco-development Myforest 22(2) 97-101

- Sugatham, R. 1981. A survey of Ceylon frog mouth (*Batrachostomus moniliger*) habitat in the western ghats of India. J. Bombay Natl. Hist. Soc. 78(2): 309-316
- Sukumar, R. 1985. Ecology and conservation of the Asian elephant in South India (with special reference to the Chamarajnagar and Satyamangala forest divisions) Tech Report No. 14 IISc., Bangalore.
- Surendra Babu, K. & Neelakantan, B. 1983. Biology of *Liza parsia* in the Kali estuary, Karwar. Mahasagar 16(3): 381-389
- Suresh, K. & Reddy, M.P.M. 1978. Distribution of nutrients in the nearshore waters off Mangalore. Mahasagar 11(3&4): 145-154
- Suresh, P.V. 1989. studies on the mangroves of the Karnataka part of the Malabar coast. Ph.D thesis submitted to Bangalore University.
- Syda Rao, G. 1988. Biology of squid *Loligo duraucelli* orbigny with a note on its fishery off Mangalore. Indian J. Fish. 35(3): 121-130
- Syda Rao, G. & Satyanarayana Rao, K. 1985. Survey of clam and oyster resources of some Karnataka estuaries. Indian J. Fish. 32: 74-89.
- Thakur, R.K. & Shivaramakrishnan, V.R. 1991. A note on the insect pest problems in Kundapur and Mangalore forests, Karnataka (South India). Myforest 27(2): 187-193
- Trimurthi, N. 1955. The forests of Coorg state. Indian For. (1)-3-6
- Ubhayankar, P.G. 1953. A note on Coastal plantations of *Casuarina equisetifolia* in North Kanara. (8): 446-451
- Untawali, A.G., Reddy, C.R.K. and Deshmukh, G.V. 1989. Ecology of intertidal benthic algae of Northern Karnataka coast. Indian J. Mar.Sci. 18: 73-83
- Uppina, S.F. 1985. Namma mangagalu. Myforest 21(2): 129-134
- Usman, S. & Puttarudraiah, M. 1955. A list of insects of Mysore including the mires. Dept. Agric. Mysore Ent. Ser. Bull No 16 pp 194
- Uttangi, J.C. 1985. Bird fauna in the urban habitats of Hubli and Dharwad. Myforest 21(3). 151-161
- Uttangi, J.C. 1993. River ecology and wildlife conservation - A review of the importance of rivers for wildlife, with special reference to avifaunal survey of forest reserves of Mahadayi valley in western ghats, India. Myforest 29(3): 179-186
- Yoganarasimhan, S.N., Mary, Z., Pattanashetty, J.K., Togunastri, V.S., Holla, B.V., Abraham, K. & Raj, P.V. 1981. New plant sources for some Ayurvedic drugs. Ind. J. For 4(2): 129-137.
- Yoganarasimhan, S.N, Govindaiah and Nair, V.K. 1982. Additions to the flora of Bangalore district. Myforest (1): 1-3
- Yoganarasimhan, S.N. & Razi, B.A. 1982. Flora of Chikmagalur district, Karnataka, India. International Book Distributors, Dehra Dun.

